



TITLE:

# PS15 An application of the geometrical phase to a biophysical problem : pump current and its fluctuation(Topological Aspects of Solid State Physics)

AUTHOR(S):

Ohkubo, Jun

---

CITATION:

Ohkubo, Jun. PS15 An application of the geometrical phase to a biophysical problem : pump current and its fluctuation(Topological Aspects of Solid State Physics). 物性研究 2009, 91(6): 699-699

ISSUE DATE:

2009-03-20

URL:

<http://hdl.handle.net/2433/142876>

RIGHT:

PS15

## An application of the geometrical phase to a biophysical problem: pump current and its fluctuation

Jun Ohkubo

ISSP, University of Tokyo

The concept of the geometrical phase is applicable for various phenomena. In this poster presentation, we will discuss a new application of the geometrical phase to a 'biophysical' problem. The problem is not quantum mechanical one, but a classical stochastic one. It has been known that a stochastic system under a periodic perturbation shows a pumping phenomenon; i.e., the periodic perturbation causes a current which is not explained by a simple average of those in the strict static cases. Recently, a current and its fluctuation in an adiabatic case (with very slow periodic perturbation) has been calculated by using the Berry phase interpretation. Because a non-adiabatic case is also important for such classical stochastic systems, we evaluate the current and its fluctuation in the non-adiabatic case. In the non-adiabatic case, the statistics of the pump current is related to the Aharonov-Anandan phase, and the Floquet theory is used to calculate it.

PS16

## Modelling the Berezinskii-Kosterlitz-Thouless Transition in the $\text{NiGa}_2\text{S}_4$

Chyh-Hong Chern

ISSP, University of Tokyo

In the two-dimensional superfluidity, the proliferation of the vortices and the anti-vortices results in a new class of phase transition, Berezinskii-Kosterlitz-Thouless (BKT) transition. This class of the phase transitions is also anticipated in the two-dimensional magnetic systems. However, its existence in the real magnetic systems still remains mysterious. Here we propose a phenomenological model to illustrate that the novel spin-freezing transition recently uncovered in the NMR experiment on the  $\text{NiGa}_2\text{S}_4$  compound is the BKT-type. The novel spin-freezing state observed in the  $\text{NiGa}_2\text{S}_4$  possesses the power-law decayed spin correlation.